arXiv:cond-mat/9911169v1 [cond-mat.stat-mech] 11 Nov 1999

Superfluid critical temperature in 3D Fermi gas with repulsion

Dmitry V. Efremov [a,1], Maxim S. Mar'enko [a], Mikhail A. Baranov [b] Maxim Yu. Kagan [a],
[a] *P.L.Kapitza Institute for Physical Problems, Ul. Kosygina 2, Moscow 117334, Russia*
[b] *Russian Research Center "Kurchatov Institute", Kurchatov Square, Moscow, 123182, Russia*

Abstract

The critical temperature of a superfluid phase transition in a Fermi gas with repulsive interaction is found. The influence of a magnetic field on the transition is analyzed. The estimates for the critical temperature for a trapped gas of ^{6}Li atoms and ^{3}He–^{4}He mixtures are presented.

Keywords: Superfluidity in neutral Fermi systems; ^{3}He-^{4}He mixtures; trapped Fermi gases

---

One of the most important questions in connection with nonconventional superconductivity is the origin of an attractive interaction. In this paper we show that a nonideal Fermi gas with purely repulsive bare interaction is unstable towards Cooper pairing with orbital momentum $l = 1$. This instability exists due to Kohn-Luttinger mechanism based on many-body effects [1].

1. Theoretical model

We consider a nonideal Fermi-gas described by the Hamiltonian:

$$\hat{H} = \sum_{\sigma=\uparrow,\downarrow}\sum_{\mathbf{p}} \xi_p \hat{a}^\dagger_{\mathbf{p}\sigma}\hat{a}_{\mathbf{p}\sigma} + g\sum_{\mathbf{p}_i}\hat{a}^\dagger_{\mathbf{p}1\uparrow}\hat{a}^\dagger_{\mathbf{p}2\downarrow}\hat{a}_{\mathbf{p}3\downarrow}\hat{a}_{\mathbf{p}4\uparrow},$$

where $\xi_p = p^2/2m - \mu$, μ is the chemical potential, g the constant of a bare point-like repulsive interparticle interaction, and different spin components, $\sigma =\uparrow,\downarrow$, are assumed to have equal masses m and concentrations n.

As it was shown in [2,3], an effective interparticle interaction originated from both the bare one and many-body effects, is attractive when two particles have a nonzero relative angular momentum l. This attractive interaction is maximal for $l = 1$ and, therefore, results in a p-wave triplet Cooper pairing with the critical temperature:

$$T_{c1} = \tilde{\varepsilon}\exp\left\{-1/\nu_F|V_1^{eff}|\right\} = \tilde{\varepsilon}\exp\left\{-12.9/\lambda^2\right\},$$

where $\nu_F = mp_F/2\pi^2$ is the density of states at the Fermi energy ($p_F = (6\pi^2 n)^{1/3}$ the Fermi momentum), $V_1^{eff} = (2\ln 2 - 1)g^2\nu_F/5$ the p-wave harmonic of the effective interaction in the Cooper channel (see [2] for details), a the scattering length ($a = mg/4\pi$ in the Born approximation), $\lambda = 2ap_F/\pi$ the gas parameter ($\lambda \ll 1$ for the considered case), and $\tilde{\varepsilon}$ the cutoff parameter of the order of the Fermi-energy, $\tilde{\varepsilon} \sim \varepsilon_F = p_F^2/2m$.

---

[1] Corresponding author. E-mail: efremov@kapitza.ras.ru



Fig. 1. T_{c1}/ε_F versus polarization α for different λ.

To fix the parameter $\tilde{\varepsilon}$ and find the critical temperature one has to keep all contributions up to λ^4 in the Bethe-Salpeter equation, that defines the critical temperature (see [5] for more details). These contributions originate from the effective interaction, retardation effects (momentum and frequency dependence of the effective interaction) and renormalization of Green functions (Z-factor and m^*).

The corresponding critical temperature, found numerically, is

$$T_{c1} \approx \frac{2}{\pi} e^C \varepsilon_F \exp\left\{ -\frac{12.9}{\lambda^2(1+4.3\lambda)} + \frac{13.4}{(1+4.3\lambda)^2} \right\},$$

where $C = 0.577$ is the Euler constant, and neglected terms are of order λ. (Note, that this formula extrapolates the expression for T_{c1} from $\lambda \ll 1$ to $\lambda < 1$.)

Magnetic field dependence of the critical temperature T_{c1} can be analyzed in the same way. We present the results for T_{c1} as a function of polarization $\alpha = (n_\uparrow - n_\downarrow)/(n_\uparrow + n_\downarrow)$ on Fig 1. It turns out, that the nonmonotonic dependence of T_{c1} on α is a result of a competition between the increase of the angular dependence of the effective interaction V^{eff} and decrease of its amplitude. (The former increases $|V_1^{eff}|$ and, hence, T_{c1}, while the later decreases.)

2. Conclusions.

In conclusion, let us mention two possible experimental applications of the presented theory. The first one is to ^{3}He-^{4}He mixtures, providing that the concentration x of ^{3}He is more than 3%. (In this case the interaction between ^{3}He atoms is repulsive.) The estimate of the critical temperature in zero magnetic field gives the value $T_{c1} \approx 5 \cdot 10^{-6}$K for the maximal concentration $x \approx 9.5\%$ (at pressure 10 bar). By applying the magnetic field this value can be increased by a factor of 6 (at polarization $\alpha \sim 40\%$), that gives a hope to observe the transition experimentally.

The second application is to trapped neutral Fermi gases. For these systems the critical temperature T_{c1} is estimated to be of the order of $10^{-7} \div 10^{-6}$K for densities $n \sim 10^{14}\text{cm}^{-3}$.

Acknowledgement

We acknowledge fruitful discussions with A.F. Andreev, H.W. Capel, I.A. Fomin, Yu. Kagan, D. Rainer, and I.M Suslov. This work was supported by INTAS grant 98-963 and RFBR grants 98-02-17077 and 97-02-16532.

References

[1] W. Kohn, J.H. Luttinger, Phys. Rev. Lett., 15, 524-526 (1965).

[2] M.Yu. Kagan, A.V. Chubukov, JETP Letters, 1988, v.47, p.525.

[3] D. Fay, A. Layzer, Phys. Rev. Lett., 20, 187-190 (1968).

[4] M.Yu. Kagan, A.V. Chubukov, JETP Letters, 1989, v.50, p.483.

[5] D.V. Efremov, M.S. Marenko, M.A. Baranov, M.Yu. Kagan, JETP, to be published.